



Veterinary Use Only

Diacalf

Effervescent Tablets
Veterinary Electrolyte



ALIVIRA

Ingredients:

Sodium chloride 2.500 mg, Potassium chloride 1.250 mg, Sodiumbicarbonate 17.000 mg, Glutamic acid 5.000 mg, Lactose monohydrate q.s.p, Citric acid 5.000 mg, Salvia Rosmarinus 200 mg, Enterococcus Faecium NCIMB 10415 $1,2 \times 10^{10}$ CFU/gr, Vanilla flavor 50 mg, Ascorbic acid 150 mg, Ponceau 4R 10 mg.

Pharmaceutical Form:

Effervescent tablet.

Product Category:

Feed additive.

Target Species:

Calf, piglet.

Product Advantages:

- Good rehydration through high levels of the important electrolytes sodium, potassium, and chloride
- A sodium bicarbonate buffer against acidosis, which neutralized pH
- Energy boost in the form of lactose
- The calves will feel better, increase energy levels and recover faster.
- Vanilla flavor provides appetite.
- E. Faecium accelerates the treatment process.

Indications:

Supportive therapy in cases of diarrhea.

Recommendation:

Calf:

- Diacalf is intended for oral administration only.
- One Diacalf tablet in 2 liters of lukewarm water (Around 37 °C) per meal. Repeat during 2 or 3 days.
- During to the next 2 days, 1 liter Diacalf solution can be mixed with 1 liter of milk or milk replacer.
- In cases of severe dehydration, Diacalf solution should be administered in 3 or 4 feeds per day.
- Normally, calves can be fed Diacalf solution exclusively for up to 4 days without harmful effects.

Pig:

- Birth to weaning: When scour occurs make Diacalf available ad libitum in a clean utensil. The amount consumed per pig depends on age.
- Weaned pig: Allow ad libitum consumption up to a maximum of 1 litre per pig per day. Provide untreated water at all times.

Packaging Size: 4x50 g

Storage:

Keep away from children. Store in a clean, dry and low humidity environment at room temperature
This is not a medicine. It is a feed additive. It is not used for treatment purposes.
Optimum storage life is 24 months from the date of production.

